



CCD Levamisole Water Soluble Powder

Composition: Levamisole hydrochloride 850mg/gm.

Appearance: white crystalline powder

Indications:

For the control of levamisole susceptible gastrointestinal roundworms of domestic poultry, pigeons, aviary pet birds in zoological collections. The most commonly encountered worms include roundworms (*Ascaridia spp.*) hairworm (*Capillaria spp.*) and caecal worm (*Heterakis spp.*).

Dosage:

The target dose rate of levamisole is 28mg/kg bodyweight, or as directed by a veterinarian. As a guide, assuming an intake of medicated water of 35ml/bird during the treatment period, mix levamisole at 800grams/ 900-1000 liters of drinking water. Alternatively, mix 8grams/10liters for smaller flocks. All medicated water should be consumed within 12 hours. Remove all other sources of water during the treatment period.

Withholding periods:

Meat: 7 days

Eggs: nil

Precautions:

Treatment is best avoided during periods of exceptionally cool or hot temperatures where water consumption will be affected.

Not for use in birds which are feeding young. Parrots appear to be less tolerant than pigeons to levamisole.

DO NOT feed to any other animal not listed above

Packaging: 120g, 500g, 1kg, 25kg

Poison Schedule: S6

For further information on this product contact
CCD Animal Health & Nutrition on 1300 666 638